

Non-Linear Derivative Interactions in Relativistic Hadrodynamics

T. Gaitanos, M. Kaskulov, U. Mosel

Institut für Theoretische Physik, Universität Giessen, Germany

The Lagrangian density of Relativistic Hadrodynamics (RHD) is extended by introducing non-linear derivative (NLD) interactions of the nucleon with the meson fields. As a consequence, the nucleon selfenergy becomes both momentum and density dependent. With a single cut-off parameter, which regulates the NLD Lagrangian, our approach is compatible with results from microscopic nuclear matter calculations as well as with Dirac phenomenology. It also reproduces the correct behavior of the real part of the proton-nucleus potential both at low and at high proton energies.

Key words: relativistic hadrodynamics, non-linear derivative model, nuclear matter, equation of state, Schrödinger equivalent optical potential

PACS numbers: 21.65.-f, 21.65.Mn, 25.40.Cm

1 Introduction

Relativistic mean-field models have been widely accepted as a successful tool for the theoretical description of different nuclear systems such as nuclear matter, finite nuclei and heavy-ion collisions [1]. The major advantage of relativistic mean-field (RMF) models has been a novel description of the saturation mechanism and an explanation of the strong spin-orbit force. An energy dependence of the Schrödinger equivalent optical potential [2] is naturally included as a consequence of a relativistic description.

The common starting basis for RMF models is the relativistic hadrodynamics (RHD) with interacting hadrons, i.e., nucleons and mesons, as the relevant degrees of freedom. Starting from the pioneering work of Duerr [3], simple Lagrangians have been introduced [4], and since then many different extensions and treatments have evolved. The Lagrangian of RHD can be treated in different approximations. For a practical application to nuclear matter and finite

nuclei one usually employs the mean-field approximation. In this approach the nucleon selfenergies become simple functions of various nucleon densities, but do not explicitly depend on energy or momentum of the nucleon. As a consequence a linear energy dependence of the optical potential arises. In principle, one can go beyond the mean-field approximation in a quantum field theoretical treatment by a systematic diagrammatic expansion. In this case, the nucleon selfenergies show a much more complicated structure that takes the non-trivial effects of the nuclear medium into account. In Dirac-Brueckner (DB) calculations of nuclear matter based on a given nucleon-nucleon interaction it is possible to extract the corresponding nucleon selfenergies [5]. They depend on both the density of the nuclear medium and the energy and momentum of the nucleon. They reproduce the empirical saturation point of nuclear matter as well as the energy dependence of the optical potential for energies up to ~ 350 MeV.

However, for practical applications the DB approach is not suitable due to its high complexity and its apparent limitations at high energies and densities. Thus, as an alternative approach to these *ab-initio* strategies for the nuclear many-body system a phenomenological treatment of the problem in the spirit of the mean-field approximation has been established. The basic Lagrangian of RHD has to be modified for a quantitative description of static nuclear systems such as nuclear matter and/or finite nuclei. Therefore, it becomes mandatory to introduce new terms, e.g., including non-linear selfinteractions of the scalar [6] and vector [7] meson fields, or to modify existing contributions in the Lagrangian, e.g., introducing density dependent meson-nucleon couplings [8,9]. The model parameters have to be fitted to properties of nuclear matter and/or atomic nuclei, since they cannot be derived in a simple manner from a microscopic description.

For a quantitative description of dynamical systems, e.g., hadron- and heavy-ion-induced reactions, the energy and momentum dependence of the nuclear mean-field becomes important. In conventional RMF models the meson-nucleon couplings are independent of the nucleon energy and momentum. This leads to a linear energy dependence of the Schrödinger equivalent optical potential with an unrealistically divergent behaviour at high energies. Indeed, analyses of proton-nucleus scattering data [2] show that the proton-nucleus optical potential levels off at proton energies of about $700 - 800$ MeV. Thus, alternative RMF approaches have been developed by including additional non-local contributions, i.e., by introducing Fock-terms, on the level of the RMF selfenergies leading to a density and energy dependence [10]. However, such a treatment is not covariant and also its numerical realization in actual transport calculations is difficult [10]. Another approach has been proposed in [11] and more recently in [12] by introducing linear order derivative couplings in the Lagrangian of RHD. These additional terms lead to an energy (or momentum) dependence of a nucleon inside the nuclear matter, and modify also the nuclear matter

equation of state (EoS). Zimanyi and Moszkowski [11] studied such gradient terms with the conclusion of a softening of the nuclear EoS. A more detailed study by Typel [12] involved the investigation of both, the density dependence of the nuclear EoS and the energy dependence of the optical potential. While *linear* derivative interactions of meson fields with nucleons explain the empirical energy dependence of the optical potential, a soft EoS results from an introduction of an explicit density dependence of the nucleon-meson couplings with additional parameters.

The purpose of the present work is to develop a covariant model, which generates both, the correct density and momentum dependence of the selfenergies in an unified framework. Furthermore, we find that the density and the momentum dependencies are indispensable and strongly correlated. The model proposed here is simple in realization and respects all the underlying symmetries of the RHD Lagrangian. The approach followed here is to extend the original Lagrangian of RHD [4] by including *non-linear* derivative interactions of meson fields with nucleons. We refer it to as non-linear derivative (NLD) model, and it is presented in the Section 2. The NLD Lagrangian (Section 2) contains not only the nucleon spinor and its first derivative, but it depends on all higher order derivatives of the nucleon field. As a consequence, the standard canonical formalism has to be generalized (Section 2.2). Field equations for nucleons and meson fields can be then derived (Section 2.3). The NLD model gives field equations with very simple structure in nuclear matter (Section 3) with density and energy dependent nucleon selfenergies. As a major advantage, both the equation of state (EoS) (density dependence) and the optical potential (energy dependence) are quantitatively well reproduced with a single parameter and the results are comparable with microscopic DB models, as in detailed discussed in Section 4. Finally, in Section 5, conclusions and an outlook complete the paper.

2 General formalism

This section describes the details of the formalism. In the first subsection the general Lagrangian density is constructed and discussed. In the second chapter various limiting cases are investigated, before the Euler-Lagrange equations for the relevant degrees of freedom are formulated.

2.1 The NLD Lagrangian

The starting point is the Lagrangian density of RHD (RHD) [4], which describes the interaction of nucleons through exchange of auxiliary meson fields

(Lorentz-scalar, σ , and the Lorentz-vector isoscalar and isovector meson fields ω^μ and $\vec{\rho}^\mu$) in the spirit of the One-Boson-Exchange (OBE) model, see for instance [5]. In conventional RHD the interaction fields couple to the spinors via the corresponding Lorentz-density operators, e.g., $\bar{\Psi}\Psi\sigma$ and $\bar{\Psi}\gamma^\mu\Psi\omega_\mu$ ($\bar{\Psi}\gamma^\mu\vec{\tau}\Psi\vec{\rho}_\mu$) for the scalar and vector sectors, respectively. In the mean-field approach to RHD the resulting mean-field potentials are only density dependent. For instance, its Lorentz-vector part rises linearly with nucleon density.

In RHD the relevant degrees of freedom are nucleons and mesons (σ , ω , ρ) characterized by the corresponding free Lagrangians and an interaction part between them:

$$\begin{aligned}\mathcal{L} = & \frac{1}{2} \left[\bar{\Psi} i \gamma_\mu \partial^\mu \Psi - (i \partial^\mu \bar{\Psi}) \gamma_\mu \Psi \right] - \bar{\Psi} \Psi m \\ & - \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ & + \frac{1}{2} m_\rho^2 \vec{\rho}_\mu \vec{\rho}^\mu - \frac{1}{4} \vec{G}_{\mu\nu} \vec{G}^{\mu\nu} + \mathcal{L}_{int} .\end{aligned}\tag{1}$$

The first line in Eq. (1) gives the (symmetrized) free Lagrangian for the nucleon field $\Psi = (\Psi_p, \Psi_n)^T$ with bare mass m . The other terms contain the standard Lagrangians for the scalar and vector mesons, σ , ω^μ and $\vec{\rho}^\mu$, with the strength tensors $F^{\mu\nu} = \partial^\mu \omega^\nu - \partial^\nu \omega^\mu$ and $\vec{G}^{\mu\nu} = \partial^\mu \vec{\rho}^\nu - \partial^\nu \vec{\rho}^\mu$ for the isoscalar and isovector meson fields, respectively.

We propose an extension of the standard RHD by introducing non-linear derivative couplings into the interaction Lagrangian density. Non-linear derivatives on the level of the Lagrangian are introduced by imposing an explicit dependence of the interaction Lagrangian density on higher order partial derivatives of the nucleon field Ψ :

$$\mathcal{L}_{int} \equiv \mathcal{L}(\Psi, \partial_\alpha \Psi, \partial_\alpha \partial_\beta \Psi, \dots, \bar{\Psi}, \partial_\alpha \bar{\Psi}, \partial_\alpha \partial_\beta \bar{\Psi}, \dots) .\tag{2}$$

with the specified (symmetrized) interaction given by:

$$\begin{aligned}\mathcal{L}_{int} = & \frac{g_\sigma}{2} \left[\bar{\Psi} \overleftarrow{\mathcal{D}} \Psi \sigma + \sigma \bar{\Psi} \overrightarrow{\mathcal{D}} \Psi \right] \\ & - \frac{g_\omega}{2} \left[\bar{\Psi} \overleftarrow{\mathcal{D}} \gamma^\mu \Psi \omega_\mu + \omega_\mu \bar{\Psi} \gamma^\mu \overrightarrow{\mathcal{D}} \Psi \right] \\ & - \frac{g_\rho}{2} \left[\bar{\Psi} \overleftarrow{\mathcal{D}} \gamma^\mu \vec{\tau} \Psi \vec{\rho}_\mu + \vec{\rho}_\mu \bar{\Psi} \vec{\tau} \gamma^\mu \overrightarrow{\mathcal{D}} \Psi \right] .\end{aligned}\tag{3}$$

The interaction between the spinor fields $\bar{\Psi}$, Ψ and the meson fields has a similar functional form as in standard RHD [4]. However, now new operators $\mathcal{D}$ acting on the nucleon fields appear, which is a generic non-linear function of

partial derivatives. The particular form of the non-linear derivative operator is not unique. Here a simple exponential form is assumed:

$$\vec{\mathcal{D}} := \exp\left(\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}\right) \quad , \quad \overleftarrow{\mathcal{D}} := \exp\left(\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}\right) . \quad (4)$$

The non-linear operators (4) explicitly depend on the partial derivate which is contracted with a dimensionless unit 4-vector v^μ . The choice of the vector field v^μ is not fixed from first principles. It can be identified with the nucleon 4-velocity u^μ :

$$v^\mu \equiv u^\mu := \frac{j^\mu}{\sqrt{j_\alpha j^\alpha}} \quad , \quad (5)$$

or with some other vector meson field:

$$v^\mu := \frac{\omega^\mu}{\sqrt{\omega_\alpha \omega^\alpha}} . \quad (6)$$

No further specification is in this context necessary. We will indeed see that rearrangement contributions generated by v^μ in the corresponding field equations cancel in nuclear matter. We thus consider it as an auxiliary unit vector. Furthermore, the additional factor $e^{m/\Lambda}$ has been introduced for convenience so that one does not have to renormalize the conventional meson-nucleon couplings $g_{\sigma,\omega,\rho}$ of the RHD Lagrangian. The non-linear derivative operator $\mathcal{D}$ contains Λ which will be interpreted as a cut-off parameter. Its value is supposed to be of natural hadronic scale of around 1 GeV.

The entire interaction part of the Lagrangian density is symmetrized in order to keep the whole Lagrangian hermitian. It is invariant under global phase transformations which guarantees baryon number conservation. Translational invariance leads to a conserved energy-momentum tensor.

2.2 Generalized formalism

Before going forward with the field equations it is useful to study particular limiting cases of the general Lagrangian (3) for the nucleons. The isovector sector is not considered here for simplicity. A series expansion is here useful for studying limiting cases ($i = \sigma, \omega, \rho$):

$$\begin{aligned}
g_i e^{\frac{-v^\beta i \vec{\partial}_\beta}{\Lambda}} &= g_i \sum_{n=0}^{\infty} \frac{(-v^\alpha i \vec{\partial}_\alpha / \Lambda)^n}{n!} \\
&= g_i - \frac{g_i}{\Lambda} v^\alpha i \vec{\partial}_\alpha + \frac{1}{2!} \frac{g_i}{\Lambda^2} v^\alpha v^\beta i \vec{\partial}_\alpha i \vec{\partial}_\beta \pm \dots .
\end{aligned}$$

The interaction terms of the Lagrangian density (3) in powers of the partial derivatives $i\partial^\mu$ takes the form

$$\mathcal{L}_{int} = \mathcal{L}_{int}^{(0)} + \mathcal{L}_{int}^{(1)} + \mathcal{L}_{int}^{(2)} + \dots . \quad (7)$$

In *zeroth order* the interaction Lagrangian is

$$\mathcal{L}_{int}^{(0)} = -\frac{1}{2} g_\omega \left(\bar{\Psi} \gamma^\mu \Psi \omega_\mu + \omega_\mu \bar{\Psi} \gamma^\mu \Psi \right) + \frac{1}{2} g_\sigma \left(\bar{\Psi} \Psi \sigma + \sigma \bar{\Psi} \Psi \right) . \quad (8)$$

Up to zeroth order in the expansion of the non-linear derivative terms the standard Lagrangian of RHD is therefore retained:

$$\mathcal{L}_{RHD} = \frac{1}{2} \left[\bar{\Psi} \gamma_\mu i D^\mu \Psi + (\bar{\Psi} i D^\mu) \gamma_\mu \Psi \right] - \bar{\Psi} \Psi (m - g_\sigma \sigma) , \quad (9)$$

with minimal meson-nucleon coupling terms and introducing the covariant derivatives $iD^\mu := i\partial^\mu - g_\omega \omega^\mu$. Our general formalism contains as a limiting case, i.e., $\Lambda \rightarrow \infty$, the conventional RHD.

The consideration of higher orders leads to modified Lagrangians. E.g., up to first order a linear dependence on these derivatives is obtained, and so forth. However, a truncation of the expansion to an arbitrary order is not obvious. Instead, the most general case of a *resummation of all higher non-linear orders* in the interaction Lagrangian is meaningful, as introduced in Eq. (3). For this reason the full formalism will be considered for the rest of this work.

The NLD Lagrangian $\mathcal{L}$ is a functional of not only $\Psi, \bar{\Psi}$ and their first derivatives, but it depends on all higher order space-time derivatives of the spinor fields Ψ and $\bar{\Psi}$. Thus the standard expressions for the Euler-Lagrange equations and the Noether Theorem do not apply. Instead of, we have to explicitly derive the generalized Noether Theorem for a Lagrangian density which contains higher order derivatives, e.g., $\mathcal{L} = \mathcal{L}(\phi, \partial_\alpha \phi, \partial_\alpha \partial_\beta \phi, \dots)$. This step is important in defining conserved quantities such as the energy-momentum tensor and Noether currents.

For a general functional containing fields and their higher order derivatives the Euler-Lagrange equations take the form:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \phi} - \partial_\alpha \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \phi)} + \partial_\alpha \partial_\beta \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \partial_\beta \phi)} + \dots + \\ (-)^n \partial_{\alpha_1} \partial_{\alpha_2} \dots \partial_{\alpha_n} \frac{\partial \mathcal{L}}{\partial (\partial^{\alpha_1} \partial^{\alpha_2} \dots \partial^{\alpha_n})} = 0 . \end{aligned} \quad (10)$$

For a general transformation of the fields and the coordinates the variational method for an arbitrary Lagrangian density (2), which contains higher order derivatives, leads to the following continuity equation (see appendix A for detailed derivations):

$$\partial_\mu [J^\mu - T^{\mu\nu} \delta x_\nu] = 0 . \quad (11)$$

where J^μ and $T^{\mu\nu}$ are defined as:

$$\begin{aligned} J^\mu = & \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} - \partial_\beta \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \phi)} + \partial_\beta \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\gamma \phi)} \mp \dots \right] \delta \phi_T \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\gamma \phi)} \pm \dots \right] \partial_\beta \delta \phi_T \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\nu \partial_\xi \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\gamma \partial_\nu \partial_\xi \phi)} \pm \dots \right] \partial_\nu \partial_\xi \delta \phi_T \\ & + \dots \end{aligned} \quad (12)$$

$$\begin{aligned} T^{\mu\nu} = & \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} - \partial_\beta \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \phi)} + \partial_\beta \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\gamma \phi)} \mp \dots \right] \partial^\nu \phi \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\gamma \phi)} \pm \dots \right] \partial_\beta \partial^\nu \phi \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\xi \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\gamma \partial_\beta \partial_\xi \phi)} \pm \dots \right] \partial_\beta \partial_\xi \partial^\nu \phi + \dots - g^{\mu\nu} \mathcal{L} . \end{aligned} \quad (13)$$

In case of Poincare' transformations, i.e., $\delta \phi_T = \phi'(x') - \phi(x) = 0$ and $x' = x^\mu + \delta x^\mu$ we obtain $\partial_\mu T^{\mu\nu} = 0$ and therefore the conserved energy-momentum tensor density. In case of global phase transformations, i.e., $\delta \phi_T \approx i\epsilon \phi$ and $\delta x^\mu = 0$, the continuity equation $\partial_\mu J^\mu = 0$ is retained.

Contrary to the standard Noether theorem, now more terms proportional to higher order derivatives of the variation of the field $\delta \phi$ appear, and for each order with respect to the derivative on $\delta \phi$, an infinite series appears. However, we will see that the infinite series can be resummed to compact expressions and that all the equations will simplify considerably.

2.3 Field equations for all degrees of freedom

The application of the generalized Euler-Lagrange equations (11) to the full Lagrangian density (3) with respect to the Spinor field Ψ leads to a Dirac equation with selfenergies, which in general contain infinite series with respect to the partial derivatives¹. As shown in Appendix B, all series can be resummed to simple exponential functions leading to the following Dirac equation:

$$[\gamma_\mu(i\partial^\mu - \Sigma^\mu) - (m - \Sigma_s)]\Psi = 0, \quad (14)$$

with Lorentz-vector and Lorentz-scalar selfenergies defined as:

$$\Sigma^\mu = g_\omega \omega^\mu e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} + \Sigma_r^\mu, \quad \Sigma_s = g_\sigma \sigma e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}}. \quad (15)$$

Both vector and scalar selfenergies Σ^μ , Σ_s show an explicit linear behavior with respect to the corresponding fields, ω^μ and σ , respectively, which looks like as in the linear Walecka model of RHD. It has been widely discussed in the literature that the linear Walecka model does not reproduce the compressibility of nuclear matter at saturation density, except if selfinteractions between the σ meson fields [4,6] and the ω fields [7] are introduced. One may thus expects the appearance of the same problem also in our formalism. On the other hand, the meson fields will contain a residual non-linear density behavior, which will be discussed below, together with the rearrangement term Σ_r^μ . Apart from the density dependence, the selfenergies contain now a non-linear dependence on the partial derivatives, which in nuclear matter are related to the 4-momentum of a moving nucleon inside the nuclear matter at rest.

Proca and Klein-Gordon equations for the meson fields are immediately derived from the Lagrangian density (3) by considering the explicit ω -dependence of the 4-vector v^μ (see again Eq. (6)):

$$\partial_\mu F^{\mu\nu} + m_\omega^2 \omega^\nu = \frac{1}{2} g_\omega \left[\bar{\Psi} e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \gamma^\nu \Psi + \bar{\Psi} \gamma^\nu e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \right] + \Omega_r^\nu, \quad (16)$$

$$\partial_\alpha \partial^\alpha \sigma + m_\sigma^2 \sigma = \frac{1}{2} g_\sigma \left[\bar{\Psi} e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \Psi + \bar{\Psi} e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \right]. \quad (17)$$

¹ In order to keep the presentation of the model and the discussions transparent, we omit the ρ -meson contributions in all the expressions. It is, however, included in the calculations for neutron matter.

The Proca equation for the ω meson field (16) shows a rather complex structure of the source term indicating a highly non-linear behavior of the ω field (the rearrangement term Ω_r^μ is given in Eq. (B.8), however, it is not relevant as discussed below). The same feature also holds for the source term of the σ -meson field, although in the latter case its structure is simpler. Note that the meson field equations for the two mesons are coupled through the non-linear exponential terms. In nuclear matter these equations will simplify considerably, however, the coupling between both meson equations will remain. This feature together with the non-linearities in the source terms will generate a residual non-linear density behavior also for the vector field, which in the standard Walecka model is linear in density.

Depending on the particular choice of the auxiliary unit vector in Eqs. (5) or (6) rearrangement terms appear in the vector part of the nucleon selfenergy (Σ_r^μ) or in the source term of the Proca equation (Ω_r^μ). An important feature is, however, that independent of the choice of v^μ the rearrangement terms exactly cancel in nuclear matter (see appendix B.1 and B.2) simplifying the equations of motion to a large extent. We, therefore, do not specify these rearrangement contributions here any further.

3 Application to infinite nuclear matter

Infinite nuclear matter is characterized by a homogeneous, isotropic and stationary system at zero temperature. Due to these symmetries the field equations simplify: the densities are constant in space-time, and only the 0-component of the currents survives. Consequently, the meson fields are constant numbers and the spatial components of the Lorentz-vector ω -field vanish. In the spirit of the mean-field approximation all meson fields are treated as classical fields implying the replacement of the source term operators in the meson field equations by their expectation values. Therefore $v^\mu = \omega^\mu / \sqrt{\omega_\alpha \omega^\alpha} = (1, \vec{0})$ or $v^\mu = j^\mu / \sqrt{j_\alpha j^\alpha} = (1, \vec{0})$, since the space-like components of the usual nucleon current $j^\mu = \langle \bar{\Psi} \gamma^\mu \Psi \rangle$ vanish in average for nuclear matter at rest.

In nuclear matter solutions of the spinor field are found with the plane wave ansatz (for simplicity isospin is neglected):

$$\Psi(s, \vec{p}) = u(s, \vec{p}) e^{-ip^\mu x_\mu}, \quad (18)$$

with s and $p^\mu = (E, \vec{p})$ being the spin and 4-momentum of the nucleon, x^μ its time-space coordinate, and u the spinor field for positive energy states.

In nuclear matter the rearrangement contributions cancel and the Dirac equa-

tion (14) maintains its form with selfenergies given by:

$$\Sigma_v = g_\omega \omega_0 e^{-\frac{E-m}{\Lambda}} \quad , \quad \Sigma_s = g_\sigma \sigma e^{-\frac{E-m}{\Lambda}} . \quad (19)$$

As one can see when the nucleons are at rest, $E = m$, the exponential factor is equal to unity and the equations are reduced to the ones from the Walecka model. The selfenergies show now a transparent behavior in energy. They contain a non-linear energy dependence, in particular, they exponentially decrease with increasing energy, as also expected from Dirac Phenomenology [2]. They are linear in the meson fields as in the linear Walecka model, however, this does not necessarily imply a linear density behavior of the mean field.

Solutions of the Dirac equation are obtained by inserting (18) and (19) into the Dirac equation (14):

$$\gamma_0 E^* \Psi = (\vec{\gamma} \cdot \vec{p} + m^*) \Psi , \quad (20)$$

with

$$E^* := E - g_\omega \omega_0 e^{-\frac{E-m}{\Lambda}} \quad , \quad m^* := m - g_\sigma \sigma e^{-\frac{E-m}{\Lambda}} . \quad (21)$$

The Dirac equation (20) has the same structure as in the free space, however, with modified effective energy E^* and effective (Dirac) mass m^* . The spatial component of the 4-momentum $\vec{p}$ remains unchanged in nuclear matter. Solutions of the Dirac equation (20) are found in the usual way:

$$u(s, \vec{p}) = N \begin{pmatrix} \phi_s \\ \frac{\vec{\sigma} \cdot \vec{p}}{E^* + m^*} \phi_s \end{pmatrix} , \quad (22)$$

with spin eigenfunctions ϕ_s and with the normalization $N = \sqrt{\frac{E^* + m^*}{2E^*}}$, which guarantees the orthogonality relation of the spinors. The following dispersion relation holds

$$E^{*2} - \vec{p}^2 = m^{*2} . \quad (23)$$

Note that, E^* and m^* depend both on density and, particular, on energy implying self consistency to obtain the dispersion relation $E = E(p)$.

In nuclear matter the NLD equations of motion for σ and ω simplify significantly to standard meson field equations:

$$m_\omega^2 \omega^0 = g_\omega \rho_0 \quad , \quad m_\sigma^2 \sigma = g_\sigma \rho_s , \quad (24)$$

with the following source terms ($\kappa = 4, 2$ for nuclear matter and neutron matter):

$$\begin{aligned}\rho_s &= \langle \bar{\Psi} e^{-\frac{E-m}{\Lambda}} \Psi \rangle = \frac{\kappa}{(2\pi)^3} \int d^3p \frac{m^*}{E^*} e^{-\frac{E-m}{\Lambda}} \Theta(p - p_F), \\ \rho_0 &= \langle \bar{\Psi} \gamma^0 e^{-\frac{E-m}{\Lambda}} \Psi \rangle = \frac{\kappa}{(2\pi)^3} \int d^3p e^{-\frac{E-m}{\Lambda}} \Theta(p - p_F) .\end{aligned}\quad (25)$$

Note that, the density ρ_0 is not related to the conserved nucleon density ρ_B . Later on, this quantity will be specified explicitly. Both vector and scalar densities ρ_0 and ρ_s differ considerably from those of the conventional Walecka model due to the dispersion relation (23) and the appearance of the exponential terms, which act as *damping factors*. These effects lead to a coupled set of equations between the meson fields and the dispersion relation. They generate a residual non-linear density dependence for both the scalar field and especially the vector ω -meson.

According to the generalized Noether-theorem (13) a conserved nucleon density and an equation of state can be derived (see appendix C). The equation

$$J^0 \equiv \rho_B = \langle \bar{\Psi} \gamma^0 \Psi \rangle + \frac{g_\omega}{\Lambda} \langle \bar{\Psi} \gamma^0 e^{-\frac{E-m}{\Lambda}} \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} e^{-\frac{E-m}{\Lambda}} \Psi \rangle \sigma \quad (26)$$

characterizes the conserved nucleon density ρ_B , from which also the relation between the Fermi momentum p_F and ρ_B is uniquely determined. The expectation values $\rho_0 = \langle \bar{\Psi} \gamma^0 e^{-\frac{E-m}{\Lambda}} \Psi \rangle$ and $\rho_s = \langle \bar{\Psi} e^{-\frac{E-m}{\Lambda}} \Psi \rangle$ have been already evaluated in (25). The expectation value $\langle \bar{\Psi} \gamma^0 \Psi \rangle = \frac{\kappa}{(2\pi)^3} (4\pi) \frac{p_F^3}{3}$ is just the usual density of the Walecka model [4]. Unlike the Walecka model, which is recovered in the limiting case of $\Lambda \rightarrow \infty$, the baryon density in the NLD approach shows a more complex structure. This results from the application of the generalized Noether-theorem to the NLD Lagrangian in nuclear matter, in which the non-linear derivative terms are coupled with the meson fields, as shown in appendix C. The extra terms in Eq. (26) reflect the dressing of the nucleons with the meson clouds. The latter start to contribute at densities above saturation, e.g., at $\rho_B = 2\rho_{sat}$ the nucleons contribute about 92% and the mesons 8% to the baryon density.

The Equation of State (EoS) is obtained from the 00-component of the energy-momentum tensor $T^{\mu\nu}$:

$$\begin{aligned}T^{00} \equiv \epsilon &= \langle \bar{\Psi} \gamma^0 E \Psi \rangle + \frac{g_\omega}{\Lambda} \langle \bar{\Psi} \gamma^0 e^{-\frac{E-m}{\Lambda}} E \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} e^{-\frac{E-m}{\Lambda}} E \Psi \rangle \sigma \\ &+ \frac{1}{2} (m_\sigma^2 \sigma^2 - m_\omega^2 \omega_0^2) ,\end{aligned}\quad (27)$$




Fig. 1. Vector (ω , upper panel) and scalar (σ , lower panel) meson fields as function of the nucleon density ρ_B . Dashed: Linear Walecka model, solid: NLD model.

with the additional expectation values:

$$\begin{aligned}\rho_s^E &= \langle \bar{\Psi} E e^{-\frac{E-m}{\Lambda}} \Psi \rangle = \frac{\kappa}{(2\pi)^3} \int d^3p \frac{m^*}{E^*} E e^{-\frac{E-m}{\Lambda}} \Theta(p - p_F), \\ \rho_0^E &= \langle \bar{\Psi} E \gamma^0 e^{-\frac{E-m}{\Lambda}} \Psi \rangle = \frac{\kappa}{(2\pi)^3} \int d^3p E e^{-\frac{E-m}{\Lambda}} \Theta(p - p_F) .\end{aligned}\quad (28)$$

4 Results

The set of coupled equations (23,24) has to be solved self consistently together with the numerical integrations for the different densities (25,28). This procedure is done as function of the Fermi momentum p_F and of the momentum p . The relation between the nucleon density ρ_B and the Fermi momentum p_F is given by Eq. (26).

In principle, the NLD model contains no parameters. The original σNN and ωNN couplings can be taken from any linear Walecka model [4], as it has been done here. The cut-off parameter Λ has to be of natural size, i.e., of typical hadronic mass scale in this problem. In the following, $\Lambda = 0.770$ GeV is chosen for a quantitative discussion of the results. Of course, different cut-off parameters $\Lambda_{\sigma,\omega}$ can be also used, which may be necessary when applying the




Fig. 2. Vector (upper panel) and scalar (lower panel) selfenergies as function of nucleon density. Dashed: Linear Walecka model, solid: NLD model, filled squares: DB model [5], filled circles: Sugahara and Toki [7].

NLD model to more complex nuclear systems such as finite nuclei. However, this is not the scope of the present work.

We start the discussion with the dependence of the isoscalar meson fields as function of the nucleon density ρ_B , see Fig. 1. In the standard linear Walecka model the ω -field shows the typical linear density behavior, while the scalar meson field saturates with increasing ρ_B due to the suppression γ -factor m^*/E^* . In the NLD model, however, a saturation of the ω -field is observed too. This result is similar to that of Ref. [7], in which selfinteractions of the ω meson field were considered. This non-linear effect on the vector meson arises from the damping factor $e^{-E/\Lambda}$, as discussed above. Non-linear effects to the scalar meson, on the other hand, are moderate due to partial compensation effects between the damping factor and the m^*/E^* term.

The selfenergies $\Sigma_{s,v}$ are proportional to the meson fields (see Eqs. (19)), however, they are dominated by the factor $e^{-E/\Lambda}$ which appears explicitly and is itself also density dependent through the in-medium dispersion relation. As can be seen in Fig. 2, the scalar part (lower panel) is thus further suppressed in the NLD model compared with the standard linear Walecka model. The density dependence of the vector field in the NLD approach differs considerably from the usual linear dependence, in particular, at nucleon densities above saturation Σ_v is strongly suppressed and tends to saturate at very high densities. The agreement between the NLD model and microscopic DB calculations




Fig. 3. Equation of state for nuclear (left panel) and pure neutron matter (right panel). Dashed: Walecka model, solid: NLD model, filled squares: DB model [5].

[5] is remarkable over a wide region in nucleon density. Another possibility of suppressing the linear density dependence of the vector field was considered in Ref. [7], in which a selfinteraction term of the ω meson was introduced in the RMF Lagrangian. As can be seen again in Fig. 2, both models are very close to each other around saturation density and below, but differ considerably for $\rho_B \geq 0.3 \text{ fm}^{-3}$.

The non-linear density dependence significantly affects the equation of state (EoS), i.e., the binding energy per nucleon as function of nucleon density. This is demonstrated in Fig. 3 for nuclear and pure neutron matter. The standard linear Walecka model leads to an EoS with high stiffness resulting in a very high value for the compression modulus. The NLD model weakens the stiffness of the EoS for nuclear and pure neutron matter to a large extent. The agreement of the NLD-EoS with the underlying DB theory is successful indicating that saturation nuclear matter properties, e.g., binding energy per nucleon and compression modulus at saturation density, are fairly well reproduced by the NLD approach. It thus turns out that standard linear Walecka models without any explicit introduction of selfinteractions of the meson fields are able to describe nuclear matter properties by the introduction of non-linear derivatives with a single cut-off parameter.

Similar effects are observed in the effective mass, as shown in the upper panel




Fig. 4. Upper panel: Nucleon density dependence of the effective (Dirac) mass m^* in units of the bare nucleon mass M . Dashed: Walecka model, solid: NLD model, filled squares: DB model [5]. Lower panel: Energy dependence of the same quantity in the NLD approach.

of Fig. 4. At saturation density a value of $m^* \approx 0.65m$ is obtained, which is again very close to DB predictions. The increase of the effective (Dirac) mass at high nucleon densities ($\rho_B \geq 0.3 \text{ fm}^{-3}$) in the NLD model reflects the attractive character of the mean-field, which is consistent with the suppression of the scalar selfenergy at high densities and is quantitatively compatible with microscopic DB calculations.

An important feature of the NLD model is the prediction of an energy (or momentum) dependence of the nuclear mean-field with only one parameter. Dirac Phenomenology on elastic proton-nucleus scattering predicts a non-linear energy dependence of the Schrödinger equivalent optical potential (see below), which cannot be reproduced in linear Walecka models nor in their extensions to non-linear meson field terms. The question arises if the NLD model can reproduce this feature with the same parameter Λ as used for the density dependence.

In the following we consider the situation of a nucleon with particular momentum p (or kinetic energy E_{kin}) relative to nuclear matter at rest. The kinetic energy for an incident free nucleon with mass m and momentum p is usually defined as $E_{kin} = \sqrt{p^2 + m^2} - m$. In the nuclear medium E_{kin} reads [12,13]

$$E_{kin} = E - m = \sqrt{p^2 + m^{*2}} + \Sigma_v - m \quad . \quad (29)$$




Fig. 5. NLD vector and scalar selfenergies (solid curves, as indicated) as function of the kinetic energy at fixed saturation density. The filled squares show DB calculations taken from [13].

As a first impression Fig. 4 (panel on the bottom) shows the energy dependence of the effective (Dirac) mass at fixed nucleon density $\rho_B = \rho_{sat} = 0.16 \text{ fm}^{-3}$. In the standard linear Walecka model m^* is independent on energy, since the scalar selfenergy is momentum independent. In the NLD model, however, the effective mass non-linearly rises with energy, and in the limit $E \rightarrow \infty$ (not shown in the figure) $m^* \rightarrow m$ is achieved, since $e^{-E/\Lambda} \rightarrow 0$. This result is in remarkable agreement with Dirac phenomenology [2].

The explicit energy dependence of the scalar and vector selfenergies in the NLD model are shown in Fig. 5 (solid curves). According to Eqs. (19) an approximately exponential decrease of $\Sigma_{s,v}$ with increasing energy can be seen, which is unambiguously fixed by the same cut-off parameter Λ as used for the density dependence. The agreement with DB results is again remarkable in an energy region up to $E_{kin} \simeq 0.4 \text{ GeV}$. The DB model describes very well saturation properties of nuclear matter. It further gives a realistic energy dependence of the selfenergies in the considered energy region by using the parameters of the underlying bare NN -potential [13]. The deviation of the fields between the NLD and DB approaches for energies above $\sim 0.4 \text{ GeV}$ will be important in describing the energy dependence of the optical potential.

The energy dependence of the nuclear mean-field is usually determined by Dirac phenomenology in elastic nucleon-nucleus scattering [2]. The essential physical object here is the Schrödinger equivalent optical potential V_{opt} , which




Fig. 6. (left) Energy dependence of the Schrödinger equivalent optical potential at density $\rho_{sat} = 0.16 \text{ fm}^{-3}$. The theoretical calculations (dashed: linear Walecka model, solid: NLD model) are compared with data from Dirac phenomenology (filled squares) [2] and with DB calculations (filled stars) [13].

serves as a convenient means to characterize the in-medium interaction of a nucleon with momentum p relative to nuclear matter at rest. It is obtained by rearranging the Dirac equation (20) into a Schrödinger-like equation for the large (upper) component of the nucleon spinor, i.e., by performing a non-relativistic reduction of the Dirac equation. This procedure applied to Eq. (20) leads to a central Schrödinger-like potential V_{opt}

$$V_{opt} = \frac{E}{m} \Sigma_v - \Sigma_s + \frac{1}{2m} (\Sigma_s^2 - \Sigma_v^2) . \quad (30)$$

It is determined by the scalar and vector selfenergies and it rises linearly with energy, if the selfenergies do not depend explicitly on momentum. This is the case of the linear Walecka model, as can be seen in Fig. 6 (dashed curve). The DB model (filled stars), on the other hand, reproduces the empirical behavior of the optical potential at low energies, since the parameters of the underlying free NN-interaction are fitted to low energy scattering data [13]. The NLD model (solid curve) with its non-linear energy dependence weakens the linear stiffness of the original Walecka model to a large extent, and the empirical energy behaviour of the optical potential can be reproduced without the introduction of any further parameters.

The strong suppression of the mean field at higher energies at saturation




Fig. 7. Energy dependence of the Schrödinger equivalent optical potential at different densities, as indicated. For each density both calculations are shown, linear Walecka model (dashed) and NLD approach (solid). The potentials at $\rho_B = 0.5\rho_{sat}$ (dotted lines), $\rho_B = \rho_{sat}$ (solid lines) and $\rho_B = 2\rho_{sat}$ (dot-dashed lines) are shifted downwards by 100 MeV, 50 MeV and 0 MeV, respectively. Horizontal lines mark the range in the energy where the potentials cross zero.

density is maintained also at higher nucleon densities, as shown in Fig. 7. For highly compressed matter, as it occurs in high energy heavy ion collisions, one thus expects a significant suppression of the repulsion of the nuclear mean field in the NLD approach. The novel result of the NLD model is thus a softening of the EoS at high densities and the Schrödinger equivalent optical potential at high energies and also high compressions. These features will be of particular interest for the theoretical description of high energy heavy-ion collisions, as they will be investigated at the compressed baryonic matter experiment at the new FAIR facility at GSI [14].

5 Conclusions and outlook

In this work the conventional linear Walecka model of relativistic hadrodynamics was extended by introducing non-linear derivative operators in the interaction Lagrangian. A generalization of standard field theoretical techniques for Lagrangians depending on all higher order derivatives of a field was necessary in deriving field equations and applying the Noether theorem in a consistent way. Starting from rather complicated field equations it was possi-

ble to reduce them to compact and simple expressions by resumming all higher order derivatives.

The model was applied in nuclear matter using a single parameter of natural hadronic mass scale. Astonishing were the non-linear density effects on the vector field without the introduction of any additional selfinteraction terms in the original Lagrangian of the linear Walecka model. These effects lead to a softening of the equation of state for nuclear and pure neutron matter at high densities. It was possible to describe quantitatively well the empirically known saturation properties. The results were also comparable to predictions of microscopic DB calculations over a wide density range. The approach lead furthermore to a momentum dependence of the selfenergies. Most noticeable was the correct reproduction of the energy dependence of the Schrödinger equivalent optical potential by utilizing the same parameter. The microscopic origin or the underlying mechanism of the desired results is not relevant at the present level of study.

In principle, one could go beyond the present study and apply the model to more complex nuclear systems, e.g., finite nuclei. In this context the present model can be further extended by introducing different cut-off parameters or different non-linear operator functions than the exponential form used in this work. Another physical object of interest would be heavy ion collisions at high energies with highest baryon compression, as they are planned at the new FAIR facility at GSI, in which the density and, in particular, the energy dependence of the nuclear mean-field play an important role. The application of the present model to heavy-ion collisions in the spirit of a covariant transport theory based on the present Lagrangian would be a great challenge for the future in studying hadronic matter under extreme conditions with the ultimative goal of exploring the equation of state at supra-normal densities.

Acknowledgments: This work is supported by BMBF.

Appendix

A Generalized Euler-Lagrange equations and Noether theorem

The procedure to derive the generalized Euler-Lagrange equations follows the canonical formalism and starts from the variation of the action. Standard steps like the integration by parts lead to the following expression:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \phi} - \partial_\alpha \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \phi)} + \partial_\alpha \partial_\beta \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \partial_\beta \phi)} + \dots + \\ (-)^n \partial_{\alpha_1} \partial_{\alpha_2} \dots \partial_{\alpha_n} \frac{\partial \mathcal{L}}{\partial (\partial^{\alpha_1} \partial^{\alpha_2} \dots \partial^{\alpha_n})} = 0 . \end{aligned} \quad (\text{A.1})$$

The derivation of the conserved energy-momentum tensor and current is more involved. The conserved energy-momentum tensor and current are obtained from a variation of the Lagrangian with respect to infinitesimal changes of the field ϕ and its argument. The total variation of the field reads

$$\delta_T \phi(x) := \phi'(x') - \phi(x) = [\phi'(x') - \phi(x')] + [\phi(x') - \phi(x)] \quad (\text{A.2})$$

which in first order reduces to the usual form

$$\delta_T \phi(x) = \delta \phi(x) + \partial_\alpha \phi(x) \delta x^\alpha \quad (\text{A.3})$$

with $\delta \phi = \phi' - \phi$ and δx^μ being the small variation of the field at fixed argument and the infinitesimal change of it, respectively.

The variation of the Lagrangian density functional thus reads

$$\begin{aligned} & \mathcal{L}[\phi'(x'), \partial_\alpha \phi'(x'), \partial_\alpha \partial_\beta \phi'(x'), \dots] - \mathcal{L}[\phi(x'), \partial_\alpha \phi(x'), \partial_\alpha \partial_\beta \phi(x'), \dots] \\ & + \mathcal{L}[\phi(x'), \partial_\alpha \phi(x'), \partial_\alpha \partial_\beta \phi(x'), \dots] - \mathcal{L}[\phi(x), \partial_\alpha \phi(x), \partial_\alpha \partial_\beta \phi(x), \dots] = 0 . \end{aligned} \quad (\text{A.4})$$

The first line in Eq. (A.4) is just the variation of $\mathcal{L}$ with respect to the fields $\phi(x), \partial_\alpha \phi, \dots$ at fixed argument, $\delta \mathcal{L}$, whereas the second line gives the variation of the Lagrangian with respect to the argument which can be evaluated up to first order. In total we obtain:

$$\delta \mathcal{L} + \partial_\alpha \mathcal{L} \delta x^\alpha = 0 \quad (\text{A.5})$$

with

$$\delta \mathcal{L} = \frac{\partial \mathcal{L}}{\partial \phi} \delta \phi + \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \phi)} \delta (\partial_\alpha \phi) + \frac{\partial \mathcal{L}}{\partial (\partial_\alpha \partial_\beta \phi)} \delta (\partial_\alpha \partial_\beta \phi) + \dots \quad (\text{A.6})$$

Inserting in the first term in Eq. (A.6) the Euler-Lagrange equations (11), we obtain after the application of the product rule following expression:

$$\begin{aligned} \delta \mathcal{L} = & \partial_\mu \left\{ \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} - \partial_\beta \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \phi)} + \partial_\beta \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\gamma \phi)} \mp \dots \right] \delta \phi \right. \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\nu \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\beta \partial_\nu \phi)} \pm \dots \right] \partial_\nu \delta \phi \\ & + \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\nu \partial_\xi \phi)} - \partial_\gamma \frac{\partial \mathcal{L}}{\partial (\partial_\mu \partial_\gamma \partial_\nu \partial_\xi \phi)} \pm \dots \right] \partial_\nu \partial_\xi \delta \phi \\ & \left. + \dots \right\} . \end{aligned} \quad (\text{A.7})$$

As next step we insert the expression A.7 into (A.5) obtaining:

$$\partial_\alpha (J^\alpha - T^{\alpha\nu} \delta x_\nu) = 0 \quad (\text{A.8})$$

which is the desired result, see also (11).

B Field equations in the NLD model

In this appendix the most important steps leading to the field equations for the spinor Ψ and the meson fields σ and ω are given. The starting basis for the derivation of a Dirac equation from the Lagrangian density (3) are the generalized Euler-Lagrange equations (11). We give the terms up to second order explicitly.

In order to show the disappearance of the rearrangement terms in nuclear matter, we choose in the derivation of the Dirac equation for the auxiliary unit vector v^μ the usual nucleon current, see Eq. (5), in which rearrangement terms only in the vector selfenergy will appear. For the derivation of the Proca equation (ω meson) $v^\mu = \omega^\mu / \sqrt{\omega_\alpha \omega^\alpha}$ will be chosen, in order to show that the disappearance of the rearrangement terms in nuclear matter is of general nature, i.e., independent of the particular choice of the auxiliary vector v^μ .

B.1 Dirac equation

For the derivation of the Dirac equation we start from the generalized Euler-Lagrange equations (A.1). We derive explicitly the first three orders in the derivative expansion:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \bar{\Psi}} + \frac{\partial \mathcal{L}}{\partial v_\mu} \frac{\partial v^\mu}{\partial \bar{\Psi}} &= \frac{1}{2} \gamma_\mu i \partial^\mu \psi - m \Psi \\ &- \frac{1}{2} g_\omega \left[\gamma_\mu \Psi \omega^\mu + \gamma_\mu e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \omega^\mu \right] + \frac{1}{2} g_\sigma \left[\Psi \sigma + e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \sigma \right] \\ &- \left[\frac{1}{2} \frac{g_\omega}{\Lambda} \left(\bar{\Psi} (i \overleftarrow{\partial}_\mu) e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \gamma^\alpha \Psi \omega_\alpha + \bar{\Psi} \gamma^\alpha e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} (-i \overrightarrow{\partial}_\mu) \Psi \omega_\alpha \right) \right. \\ &\quad \left. - \frac{1}{2} \frac{g_\sigma}{\Lambda} \left(\bar{\Psi} (i \overleftarrow{\partial}_\mu) e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \Psi \sigma + \bar{\Psi} e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} (-i \overrightarrow{\partial}_\mu) \Psi \sigma \right) \right] \\ &\times \left[\frac{\gamma^\mu \Psi}{\sqrt{j_\alpha j^\alpha}} - \frac{j^\mu j_\nu \gamma^\nu \Psi}{\sqrt{j_\alpha j^\alpha}^3} \right] \end{aligned}$$

$$\begin{aligned}
\frac{\partial \mathcal{L}}{\partial(\partial_\mu \bar{\Psi})} &= -\frac{1}{2}i\gamma^\mu \psi - \frac{1}{2}\frac{g_\omega}{\Lambda}v^\mu i\gamma_\alpha \Psi \omega^\alpha + \frac{1}{2}\frac{g_\sigma}{\Lambda}v^\mu i\Psi \sigma \\
\frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\nu \bar{\Psi})} &= -\frac{1}{2}\frac{g_\omega}{(\Lambda)^2}\frac{1}{2!}v^\mu v^\nu i i\gamma_\alpha \Psi \omega^\alpha + \frac{1}{2}\frac{g_\sigma}{(\Lambda)^2}\frac{1}{2!}v^\mu v^\nu i i\Psi \sigma .
\end{aligned} \tag{B.1}$$

Inserting (B.1) into (A.1) we obtain after some trivial algebra (up to derivatives of the meson fields which in any case vanish in nuclear matter):

$$\begin{aligned}
&(\gamma_\mu i\partial^\mu - m)\Psi - \frac{1}{2}g_\omega \gamma_\mu e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \omega^\mu + \frac{1}{2}g_\sigma e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} \Psi \sigma \\
&- \frac{1}{2}g_\omega \gamma_\mu \left(1 - \frac{1}{\Lambda}v_\alpha i\partial^\alpha + \frac{1}{2!}\frac{1}{\Lambda^2}v_\alpha v_\beta i\partial^\alpha i\partial^\beta \mp \dots\right) \Psi \omega^\mu \\
&+ \frac{1}{2}g_\sigma \left(1 - \frac{1}{\Lambda}v_\alpha i\partial^\alpha + \frac{1}{2!}\frac{1}{\Lambda^2}v_\alpha v_\beta i\partial^\alpha i\partial^\beta \mp \dots\right) \Psi \sigma + \Sigma_r \Psi = 0 .
\end{aligned} \tag{B.2}$$

with the rearrangement contribution:

$$\begin{aligned}
\Sigma_r &= \left[-\frac{1}{2}\frac{g_\omega}{\Lambda} \left(\bar{\Psi}(i\overleftarrow{\partial}_\mu) e^{\frac{i\overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \gamma^\delta \Psi \omega_\delta + \bar{\Psi} \gamma^\delta e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} (-i\overrightarrow{\partial}_\mu) \Psi \omega_\delta \right) \right. \\
&+ \frac{1}{2}\frac{g_\sigma}{\Lambda} \left(\bar{\Psi}(i\overleftarrow{\partial}_\mu) e^{\frac{i\overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \Psi \sigma + \bar{\Psi} e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}} (-i\overrightarrow{\partial}_\mu) \Psi \sigma \right) \Big] \\
&\times \left[\frac{\gamma^\mu}{\sqrt{j_\alpha j^\alpha}} - \frac{j^\mu j_\nu \gamma^\nu}{\sqrt{j_\alpha j^\alpha}^3} \right] .
\end{aligned} \tag{B.3}$$

Note that, the derivatives inside the brackets act to the spinors on the right. The terms inside the brackets represent Taylor expansions which can be resumed to the exponential function $e^{\frac{-v^\beta i \vec{\partial}_\beta + m}{\Lambda}}$. This leads us to the following Dirac equation

$$[\gamma_\mu(i\partial^\mu - \Sigma^\mu) - (m - \Sigma_s)]\Psi = 0 , \tag{B.4}$$

as it was written in Eq. (14).

In infinite nuclear matter only the 0-component of the ω meson field survives, and in the mean-field approximation all density operators are replaced by expectation values. Furthermore only the 0-component of the usual nucleon current $j^\mu = \langle \bar{\Psi} \gamma^\mu \Psi \rangle$ remains. Thus, the rearrangement contribution reads:

$$\Sigma_r = \left[\frac{g_\omega}{\Lambda} \langle \bar{\Psi} p_\mu e^{-\frac{E-m}{\Lambda}} \gamma^0 \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} p_\mu e^{-\frac{E-m}{\Lambda}} \Psi \rangle \sigma \right] \left[\frac{\gamma^\mu}{\sqrt{j_\alpha j^\alpha}} - \frac{j^\mu j_\nu \gamma^\nu}{\sqrt{j_\alpha j^\alpha}^3} \right] , \tag{B.5}$$

and by evaluating the contraction between the expressions inside the two brackets we obtain:

$$\begin{aligned} \Sigma_r = & \left[\frac{g_\omega}{\Lambda} \langle \bar{\Psi} p_0 e^{-\frac{E-m}{\Lambda}} \gamma^0 \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} p_0 e^{-\frac{E-m}{\Lambda}} \Psi \rangle \sigma \right] \left[\frac{\gamma^0}{j^0} - \frac{\gamma^0}{j^0} \right] \\ & - \left[\frac{g_\omega}{\Lambda} \langle \bar{\Psi} \vec{p} e^{-\frac{E-m}{\Lambda}} \gamma^0 \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} \vec{p} e^{-\frac{E-m}{\Lambda}} \Psi \rangle \sigma \right] \frac{\vec{\gamma}}{j^0} = 0 . \end{aligned} \quad (\text{B.6})$$

The first term in Eq. (B.6) is trivially zero. In the second line of Eq. (B.6) both expectation values are odd functions of the vector $\vec{p}$ and therefore they vanish.

B.2 Meson field equations of motion

For the meson fields σ and ω the derivations are straightforward, since here one has to use the standard Euler-Lagrange equations. The following Proca equation which includes the rearrangement and the source terms is readily derived:

$$\partial_\mu F^{\mu\nu} + m_\omega^2 \omega^\nu = \frac{1}{2} g_\omega \left[\bar{\Psi} e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \gamma^\nu \Psi + \bar{\Psi} \gamma^\nu e^{\frac{-v^\beta i \overrightarrow{\partial}_\beta + m}{\Lambda}} \Psi \right] + \Omega^\nu \quad (\text{B.7})$$

The rearrangement term in the Proca equation appears if for the auxially vector $v^\mu = \omega^\mu / \sqrt{\omega_\alpha \omega^\alpha}$ is chosen. It takes the following form:

$$\begin{aligned} \Omega^\nu := & -\frac{1}{2} \frac{g_\omega}{\Lambda} \left[\bar{\Psi} e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \left(\frac{i \overleftarrow{\partial}^\nu}{\sqrt{\omega_\alpha \omega^\alpha}} - \frac{i \overleftarrow{\partial}^\alpha \omega_\alpha \omega^\nu}{\sqrt{\omega_\alpha \omega^\alpha}^3} \right) \gamma^\delta \Psi \omega_\delta \right. \\ & \left. + \bar{\Psi} \left(-\frac{i \overrightarrow{\partial}^\nu}{\sqrt{\omega_\alpha \omega^\alpha}} + \frac{\omega_\alpha \omega^\nu i \overrightarrow{\partial}^\alpha}{\sqrt{\omega_\alpha \omega^\alpha}^3} \right) e^{\frac{-v^\beta i \overrightarrow{\partial}_\beta + m}{\Lambda}} \gamma^\delta \Psi \omega_\delta \right] \\ & + \frac{1}{2} \frac{g_\sigma}{\Lambda} \left[\bar{\Psi} e^{\frac{i \overleftarrow{\partial}_\beta v^\beta + m}{\Lambda}} \left(\frac{i \overleftarrow{\partial}^\nu}{\sqrt{\omega_\alpha \omega^\alpha}} - \frac{i \overleftarrow{\partial}^\alpha \omega_\alpha \omega^\nu}{\sqrt{\omega_\alpha \omega^\alpha}^3} \right) \Psi \sigma \right. \\ & \left. + \bar{\Psi} \left(-\frac{i \overrightarrow{\partial}^\nu}{\sqrt{\omega_\alpha \omega^\alpha}} + \frac{\omega_\alpha \omega^\nu i \overrightarrow{\partial}^\alpha}{\sqrt{\omega_\alpha \omega^\alpha}^3} \right) e^{\frac{-v^\beta i \overrightarrow{\partial}_\beta + m}{\Lambda}} \Psi \sigma \right] \end{aligned} \quad (\text{B.8})$$

In nuclear matter and in the mean-field approximation the rearrangement contribution Ω^μ to the vector meson vanishes for the same reasons as for that rearrangement term for the nucleon field (see previous subsection).

C Derivation of the Noether current

Here the most important steps leading to the Noether current (26) starting from the general expression (12) are explicitly shown. The derivation of the energy-momentum tensor (13) is then straightforward. We show that all higher order derivative terms can be resummed to simple exponential expressions for nuclear matter at rest in the mean-field approximation.

The derivative of the Lagrangian density (3) with respect to higher order partial derivatives has been demonstrated in the appendix B.1, see in particular Eq. (B.1), thus we don't repeat this intermediate step.

We consider the Noether current in the NLD model starting from the general expression (12) and taking only terms up to third order. With $\delta\bar{\Psi} = -i\epsilon\bar{\Psi}$ and $\delta\Psi = i\epsilon\Psi$ we obtain:

$$\begin{aligned}
\frac{J^\mu}{i\epsilon} = & \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\Psi)} - \partial_\beta \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\Psi)} + \partial_\beta\partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\partial_\gamma\Psi)} \right] \Psi \\
& + \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\Psi)} - \partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\partial_\gamma\Psi)} \right] \partial_\beta\Psi \\
& + \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\nu\partial_\xi\Psi)} - \partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\gamma\partial_\nu\partial_\xi\Psi)} \right] \partial_\nu\partial_\xi\Psi \\
& - \bar{\Psi} \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\bar{\Psi})} - \partial_\beta \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\bar{\Psi})} + \partial_\beta\partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\partial_\gamma\bar{\Psi})} \right] \\
& + \partial_\beta\bar{\Psi} \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\bar{\Psi})} - \partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\beta\partial_\gamma\bar{\Psi})} \right] \\
& + \partial_\nu\partial_\xi\bar{\Psi} \left[\frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\nu\partial_\xi\bar{\Psi})} - \partial_\gamma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\partial_\gamma\partial_\nu\partial_\xi\bar{\Psi})} \right] . \tag{C.1}
\end{aligned}$$

We rewrite now Eq. (C.1) by separating the terms between the different orders in the partial derivatives:

$$\frac{J^\mu}{i\epsilon} = \mathcal{O}^{(0)} + \mathcal{O}^{(1)} + \mathcal{O}^{(2)} + \dots + \mathcal{O}^{(n)} \tag{C.2}$$

with the different contributions given by:

$$\mathcal{O}^{(0)} = \left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\Psi)}\Psi - \bar{\Psi}\frac{\partial\mathcal{L}}{\partial(\partial_\mu\bar{\Psi})} \right) , \tag{C.3}$$

$$\begin{aligned}\mathcal{O}^{(1)} = & - \left(\partial_\beta \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \Psi)} \Psi - \bar{\Psi} \partial_\beta \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \bar{\Psi})} \right) \\ & + \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \Psi)} \partial_\beta \Psi - \partial_\beta \bar{\Psi} \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \bar{\Psi})} \right),\end{aligned}\quad (\text{C.4})$$

$$\begin{aligned}\mathcal{O}^{(2)} = & + \left(\partial_\beta \partial_\gamma \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \Psi)} \Psi - \bar{\Psi} \partial_\beta \partial_\gamma \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \bar{\Psi})} \right) \\ & - \left(\partial_\gamma \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \Psi)} \partial_\beta \Psi - \partial_\beta \bar{\Psi} \partial_\gamma \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \bar{\Psi})} \right) \\ & + \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \Psi)} \partial_\beta \partial_\gamma \Psi - \partial_\beta \partial_\gamma \bar{\Psi} \frac{\partial \mathcal{L}}{\partial(\partial_\mu \partial_\beta \partial_\gamma \bar{\Psi})} \right).\end{aligned}\quad (\text{C.5})$$

In mean-field approximation it turns out that first order terms (in energy) appear twice, second order terms appear three times, and so forth:

$$\mathcal{O}^{(0)} = \langle \bar{\Psi} \gamma^0 \Psi \rangle + \frac{g_\omega}{\Lambda} \langle \bar{\Psi} \gamma^0 \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \langle \bar{\Psi} \Psi \rangle \sigma, \quad (\text{C.6})$$

$$\mathcal{O}^{(1)} = -\frac{g_\omega}{\Lambda} \frac{2}{2!} \langle \bar{\Psi} \gamma^0 \frac{E}{\Lambda} \Psi \rangle \omega_0 + \frac{g_\sigma}{\Lambda} \frac{2}{2!} \langle \bar{\Psi} \frac{E}{\Lambda} \Psi \rangle \sigma, \quad (\text{C.7})$$

$$\mathcal{O}^{(2)} = \frac{g_\omega}{\Lambda} \frac{3}{3!} \langle \bar{\Psi} \gamma^0 \frac{E^2}{\Lambda^2} \Psi \rangle \omega_0 - \frac{g_\sigma}{\Lambda} \frac{3}{3!} \langle \bar{\Psi} \frac{E^2}{\Lambda^2} \Psi \rangle \sigma, \quad (\text{C.8})$$

and for the $n - th$ -order one obtains

$$\mathcal{O}^{(n-1)} = (-)^n \frac{g_\omega}{\Lambda} \frac{n}{n!} \langle \bar{\Psi} \gamma^0 \frac{E^{n-1}}{\Lambda^{n-1}} \Psi \rangle \omega_0 - (-)^n \frac{g_\sigma}{\Lambda} \frac{n}{n!} \langle \bar{\Psi} \frac{E^{n-1}}{\Lambda^{n-1}} \Psi \rangle \sigma. \quad (\text{C.9})$$

Therefore all the infinite series of terms can be resummed into simple expressions. Adding all these terms and considering the fact that $\frac{n}{n!} = \frac{1}{(n-1)!}$ we obtain the desired result, see Eq. (26).

References

- [1] B. Serot and J.D. Walecka, Int. J. Mod. Phys. **E6** (1997) 631.
- [2] E.D. Cooper, et al., Phys. Rev. **C47** (1993) 297.
- [3] H.P. Duerr, Phys. Rev. **103** (1956) 469.
- [4] J.D. Walecka, Ann. Phys. (NY) **83** (1974) 491.
- [5] R. Brockmann and R. Machleidt, Phys. Rev. **C42** (1990) 1965.
- [6] J. Boguta, A.R. Bodmer, Nucl. Phys. **A292** (1977) 413.
- [7] Y. Sugahara and H. Toki, Nucl. Phys. **A579** (1994) 557.
- [8] R. Brockmann and H. Toki, Phys. Rev. Lett. **68** (1992) 3408.
- [9] C. Fuchs, H. Lenske and H.H. Wolter, Phys. Rev. **C52** (1995) 3043.
- [10] B. Blättel, V. Koch and U. Mosel, Rep. Prog. Phys. **56** (1993) 1.
- [11] J. Zimanyi and S.A. Moszkowski, Phys. Rev. **C42** (1990) 1416.
- [12] S. Typel, Phys. Rev. **C71** (2005) 064301.
- [13] B. ter Haar and R. Malfliet, Phys. Rev. **C36** (1987) 1611;
B. ter Haar and R. Malfliet, Phys. Rep. **149** (1987) 207.
- [14] P. Senger, Phys. Part. Nucl. **39** (2008) 1055.